

The Antarctic Tern is not easily distinguished from some Palaearctic terns, and hitherto it has often been overlooked in South Africa. A specimen collected at Cape St. Francis on 23 July 1936 and stated to be a Common Tern, *S. hirundo*, (illustrated in Hewitt, 1937) has been identified as a juvenile Antarctic Tern (Liversidge, 1957).

Another early record, by Courtenay-Latimer (1957), is confusing and incomplete, and her report of the Antarctic Tern breeding in South Africa on Stag Island, Algoa Bay, in the winter of 1940 has been accepted by modern texts (McLachlan & Liversidge, 1970; Watson, 1975). The breeding record is open to doubt, since, apart from the apparently abnormal breeding season, adult Antarctic Terns moult while in South Africa (pers. obs.) and it is unusual for terns to breed while moulting. Elsewhere in its range the bird is strictly a summer breeder (Berruti & Harris, 1976; Hagen, 1952; Parmalee & Maxson, 1974) The record has never been subsequently verified.

#### Acknowledgements

I thank K. Z. Edwards, P. G. H. Frost, P. A. R. Hockey, J. S. Pringle, W. R. Siegfried, J. C. Sinclair and R. W. Summers for observations of Antarctic Terns in South Africa. The Projects Fund of the South African Ornithological Society gave financial assistance to J. S. Pringle, enabling seasonal counts to be made at Kommetjie. I acknowledge

financial and other support from the Sea Fisheries Branch, the South African Nature Foundation and the South African National Foundation for the Conservation of Coastal Birds, during 1971 and 1972, and the University of Cape Town for financial support during 1975 and 1976.

#### References

- Berruti, A. & Harris, A. Breeding schedules of Antarctic and Kerguelen Terns at Marion Island. *Notornis*, **23**, 39-41, 1976.
- Courtenay-Latimer, M. On the breeding of *Sterna vittata tristanensis* Murphy off the coast of Cape Province, South Africa. *Bull. Brit. ornithol. Club*, **77**, 82-83, 1957.
- Elliott, H.F.I. A contribution to the ornithology of the Tristan da Cunha Group. *Ibis*, **99**, 545-586, 1957.
- Hagen, Y. Birds of Tristan da Cunha. *Results of the Norwegian Scientific Expedition to Tristan da Cunha 1937-1938*, **20**, 1-248, 1952.
- Hewitt, J. Several new and noteworthy records of sea-birds from the eastern coast of the Cape Province. *Ostrich*, **8**, 1-9, 1937.
- Liversidge, R. On the status of the Tristan Tern in South Africa. *Ostrich*, **28**, 176-177, 1957.
- McLachlan, G.R. & Liversidge, R. *Roberts Birds of South Africa*. Cape Town, John Voelker Bird Book Fund, 1970.
- Parmalee, D.F. & Maxson, S.J. The Antarctic Terns of Anvers Island. *Living Bird*, **13**, 233-250, 1974.
- Watson, G.E. *Birds of the Antarctic and Sub-Antarctic*. Washington, American Geophysical Union, 1975.
- (Received October 1976).

## A note on the daily variation of the geomagnetic vertical intensity at Marion Island

P.R. Sutcliffe

Antarctic Geomagnetic Programme,  
Magnetic Observatory of the CSIR, Hermanus.

*Significant phase shifts occur in the daily variation of the geomagnetic vertical intensity at Marion Island. These phase shifts are not due to the 'island effect', as previously reported, but are brought about by the  $L_2(Z)$  variation and are functions of the lunar phase angle.*

*Beduidende faseverskuiwings kom voor in die daaglikse variasie van die geomagnetiese vertikale intensiteit op Marion Eiland. Dié faseverskuiwings is nie te wyte aan die 'eiland-effek' soos voorheen vermeld nie, maar word teweeggebring deur die  $L_2(Z)$  variasie en is 'n funksie van die maan se fasehoek.*

### Introduction

The daily variation of the geomagnetic vertical intensity  $Z$  at inland observatories on quiet days consists essentially of the solar quiet day variation  $Sq(Z)$  plus a small perturbation due to the lunar daily variation  $L(Z)$ . However, the daily variation at coastal and oceanic island observatories is sometimes significantly modified by other factors. Malin (1969) reported an anomalously large value of the lunar semi-diurnal variation  $L_2(Z)$  at coastal observatories and ascribed it to the generation of electric currents in the sea due to tidal movements of the conductive water across the geomagnetic field. The anomalous behaviour of the magnetic vertical intensity on oceanic islands, known as the 'island effect', has been reported by Mason (1963), Voppel (1964), and Sasai (1967).

Since an island acts as a region of low conductivity in a sheet of high conductivity, currents induced in the ocean are compelled to flow around the island; consequently the vertical component of the magnetic field produced by the induced

current will be in opposite senses on opposite sides of the island. Numerous workers (Mason, 1963; Voppel, 1964; Sasai, 1967) have reported a complete reversal in sign for short-period disturbances of one hour or less, while Mason (1963) found phase shifts of up to  $70^\circ$  in the daily variation. Sasai (1967) concluded that the island effect, which is observed for the short-period range, vanishes at a period ranging from 8 to 24 hours. Rikitake (1970) confirmed theoretically that the phase shift is a function of the frequency of the variation. Kühn and Sutcliffe (1972) however, presented evidence of a phase reversal in  $Sq(Z)$  at Marion Island ( $46^\circ 52' S, 37^\circ 50' E$ ) and attributed it to the 'island effect'. In this note we show that substantial phase shifts are observed in the daily variation of  $Z$  at Marion, but that these can be explained without requiring the inconsistency of the frequency dependence of the island effect as suggested by Kühn and Sutcliffe (1972).

### Data selection and analysis

The observations on which Kühn and Sutcliffe (1972) based their conclusions were made with a BMZ on 10 and 11 May 1971. Subsequently, a magnetic observatory was established on Marion Island and hourly mean values for the period 1 June 1973-31 May 1975 are available. Sutcliffe (1977) utilized these data to study  $Sq$  at Marion. He found that the range of the mean  $Sq(Z)$  is anomalously small, especially during equinoctial and winter months, but found no evidence of a phase reversal.

Provided there is no significant magnetic activity present, the daily variation observed on any specific day will consist of  $Sq$  modified in a regular way, with a period of half a lunar




Fig. 2. Harmonic dials for the semi-diurnal waves of the daily variation of  $Z$  during the 12 lunar phases. Vector end-points are represented by circles and numbered according to lunar phase angle. The triangle indicates the end-point of the  $Sq_2(Z)$  vector.

$Z$  on 10 and 11 May 1971. According to Equation 1 the lunar phase angles were  $\nu = 0$  and  $\nu = 1$  respectively on these two days. In Fig. 3 the following curves, displaced vertically with respect to each other, are plotted:

(a) the mean synthesised daily variation for days on which  $\nu = 0$  and  $\nu = 1$ ,




Fig. 3. The curves represent: (a) the mean daily variation at Marion when  $\nu = 0$  and  $\nu = 1$ , (b) the mean variation at Marion on 10 and 11 May 1971, (c) the mean  $Sq(Z)$  at Marion Island, and (d) the mean  $Sq(Z)$  at Hermanus for the months April, May and June.

- (b) the mean variation determined from the BMZ observations on 10 and 11 May 1971,  
 (c) the mean  $Sq(Z)$  at Marion for the months April, May and June, and  
 (d) the mean  $Sq(Z)$  at Hermanus for the months April, May and June.

Curve (d) is similar to the winter solstice  $Sq(Z)$  curve for dip-latitude  $50^\circ S$  (Matsushita, 1967) which was used by Kühn and Sutcliffe (1972) for comparison with curve (b). The difference between curves (b) and (d) is clear and led them to conclude, albeit erroneously, that there was a phase reversal of  $Sq(Z)$  at Marion. From Fig. 1b we see that during daylight hours the semi-diurnal variation, on days on which  $\nu = 0$  and  $\nu = 1$ , attains its maximum value at approximately 0930 LT, while the maximum in  $Sq_2(Z)$  occurs at 1230 LT; the effect this has on the daily variation is illustrated by the difference between curves (a) and (b) in Fig. 3, together with the conclusions drawn from Fig. 1b, demonstrate that the apparent phase shift in  $Sq(Z)$  on 10 and 11 May 1971 was due to the small  $Sq(Z)$  variation being modified by  $L_2(Z)$ .

## Conclusions

There is no evidence of a significant phase shift in  $Sq(Z)$  on the north-east coast of Marion Island as a result of the island effect. Nevertheless, significant phase shifts occur in the daily variation of the geomagnetic vertical intensity; these are brought about by the modification of  $Sq(Z)$  by  $L_2(Z)$ . The shape of the daily variation is thus a function of lunar phase angle and will very nearly repeat itself twice per lunation.

The conclusions drawn in this paper are based on data for a very limited period. A detailed study of the lunar daily variation at Marion Island will be made when more data become available.

## Acknowledgements

I thank G. L. M. Scheepers for discussion. The Geomagnetic Programme at Marion Island is financially and logistically supported by the South African Department of Transport.

## References

- Chapman, S. and Bartels, J. *Geomagnetism*. Oxford, Clarendon Press, 1940.  
 Kühn, G.J. and Sutcliffe, P.R. Evidence of a phase reversal in the quiet day variation of the magnetic vertical intensity at Marion Island. *S. Afr. J. Antarct. Res.*, **2**, 45-48, 1972.  
 Malin, S.R.C. The effect of the sea on lunar variations of the vertical component of the geomagnetic field. *Planet. Space Sci.*, **17**, 487-490, 1969.  
 Mason, R.G. Spatial dependence of time-variations of the geomagnetic field in the range 24 hrs.- 3 mins. on Christmas Island. Ref. 63-3, Geophysics Dept., Imperial College of Science and Technology, London, 1963.  
 Matsushita, S. Solar quiet and lunar daily variation fields. In *Physics of Geomagnetic Phenomena I*, edited by S. Matsushita and W.H. Campbell. New York, Academic Press, 1967.  
 Parkinson, W.D. An analysis of the geomagnetic diurnal variation during the International Geophysical Year. *Gerlands Beitr. Geophysik*, **80**, 199-232, 1971.  
 Rikitake, T. A theory of frequency dependent effect of an Island on geomagnetic variations. *Bull. Earthquake Res. Inst. Tokyo Univ.*, **48**, 189-203, 1970.  
 Sasai, Y. Spatial dependence of short-period geomagnetic fluctuations on Oshima Island. *Bull. Earthquake Res. Inst. Tokyo Univ.*, **45**, 137-157, 1967.  
 Schlapp, D.M. World-wide morphology of day-to-day variability of  $Sq$ . *J. atmos. terr. Phys.*, **30**, 1761-1776, 1968.  
 Sutcliffe, P.R. The geomagnetic  $Sq$  variation at Marion Island. *S. Afr. J. Sci.*, **73**, 173-178, 1977.  
 Voppel, D. Erdmagnetische Variationen auf ozeanischen Inseln. *Dt. hydrogr. Z.*, **17**, 179-182, 1964.